Scored Patient-Generated Subjective Global Assessment (PG-SGA)

| Scored Patient-Generated Subjective | Patient ID Information | | | | | | | |
|--|---|--|--|--|--|--|--|--|
| Global Assessment (PG-SGA) | Robinson, James | | | | | | | |
| History (Boxes 1-4 are designed to be completed by the patient.) | | | | | | | | |
| Weight (See Worksheet 1) In summary of my current and recent weight: | 2. Food Intake: As compared to my normal intake, I would rate my food intake during the past month as: unchanged (9) | | | | | | | |
| I currently weigh about 150 pounds I am about 5 feet 10" tall | more than usual (0) peless than usual (1) I am now taking normal food but less than normal amount (1) | | | | | | | |
| One month ago I weighed aboutpounds Six months ago I weighed aboutpounds During the past two weeks my weight hes: | ☐ little solid food (2) ☐ only liquids (3) ☐ only nutritonal supp lements (3) ☐ very little of anything | | | | | | | |
| During the past two weeks my weight has: Decent of the past two weeks my weight has: Decent of the past two weeks my weight has: Decent of the past two weeks my weight has: Decent of the past two weeks my weight has: Decent of the past two weeks my weight has: Decent of the past two weeks my weight has: Decent of the past two weeks my weight has: Decent of the past two weeks my weight has: Decent of the past two weeks my weight has: Decent of the past two weeks my weight has: Decent of the past two weeks my weight has: Decent of the past two weeks my weight has: Decent of the past two weeks my weight has: Decent of the past two weeks my weight has: Decent of the past two weeks my weight has: Decent of the past two weight has: Dece | very little of anything (4) only tube feedings or only nutrition by vein (9) Box 2 2 | | | | | | | |
| 3. Symptoms: I have had the following problems that have kept me from eating enough during the past two weeks (check all | 4. Activities and Function: Over the past month, I would generally rate my activity as: | | | | | | | |
| that apply): no problems eating (0) | normal with no limitations (0) | | | | | | | |
| no appenite, just did not feel like eating | not my normal self, but able to be up and | | | | | | | |
| nausea (i) vomiting (3) | about with fairly normal activities (1) | | | | | | | |
| constipation (1) diarrhea (3) | not feeling up to most things, but in bed or chair | | | | | | | |
| mouth sores (2) | less than half the day (2) | | | | | | | |
| things taste funny or have no taste (1) smells bother me (1) | able to do litt le activity and spend most | | | | | | | |
| problems swallowing (2) feel full quick ly (1) | of the day in bed or chair (3) | | | | | | | |
| pain; where? (1) | pretty much bedridden, rarely out of bed Box 4 3 | | | | | | | |
| Examples: depression, money, or dental problems Box 3 | Additive Score of the Boxes 1-4 14 A | | | | | | | |
| The remainder of this form will be completed by your doctor, nurse, or therapist. Thank you. | | | | | | | | |
| 5. Disease and its relation to nutritional requirements (See Work | sheet 2) | | | | | | | |
| All relevant diagnoses (specify) | | | | | | | | |
| Primary disease stage (circle if known or appropriate) I II I. Age 72 | II (V Other | | | | | | | |
| 6. Metabolic Demand (See Worksheet 3) | | | | | | | | |
| 6. Metabone Delinand (Dec 1701 Girect 3) | Numerical score from Worksheet 3 C | | | | | | | |
| 7. Physical (See Worksheet 4) | Numerical score from Worksheet 4 D | | | | | | | |
| Global Assessment (See Worksheet 5) | Total PG-SGA score | | | | | | | |
| ☐ Well-nourished or anabolic (SGA-A) Moderate or suspected malnutrition (SGA-B) ☐ Severely malnourished (SGA-C) | (Total numerical score of A+B+C+D above) (See triage recommendations below) | | | | | | | |
| Clinician Signature Churche Church RD BN PA MD DO Orber Date 76/06 | | | | | | | | |
| Nutritional Triage Recommendations: Additive score is used to define specific nutritional interventions including patient & family education, symptom management including pharmacologic intervention, and appropriate nutrient intervention (food, nutritional supplements, enteral, or parenteral triage). First line nutrition intervention includes o ptimal symptom management. No intervention required at this time. Re-assessment on routine and regular basis during treatment. Patient & family education by dietitian, nurse, or other clinician with pharmacologic intervention as indicated by symptom survey (Box 3) and laboratory values as appropriate. Requires intervention by dietitian, in conjunction with nurse or physician as indicated by symptoms survey (Box 3). Indicates a critical need for improved symptom management and/or nutrient intervention options. | | | | | | | | |

Global PG-SGA rating (A, B, or C) = 3

Worksheets for PG-SGA Scoring

Do FD Onery
Boxes 1-4 of the PG-SGA are designed to be completed by the patient. The PG-SGA numerical score is determined using
the parenthetical points noted in boxes 1-4 and 2) the worksheets below for items not marked with parenthetical points. Scores for

| boxes 1 and 3 are addit | ive within each box | and scor | es for box | ces 2 and | 4 are based on the high | est scored iter | n checker | off by | the patient. | |
|---|---|---------------------------------|-----------------------------|--|---|-------------------|--------------------------|------------|---|--|
| Worksheet 1 - Scot To determine score, use 1 data only if there is no 1 r | month weight data if month weight data. U | evailable. se points b | Use 6 mo elow to so | core | Worksheet 2 - Sco Score is derived by addin that pertain to the patient | g I point for eac | | | | |
| weight change and add one extra point if patient has lost weight 2 weeks. Enter total point score in Box 1 of the PG-SG | | | | ng the | Category | | | Points | | |
| past 2 weeks. Enter total | bom: scote ni pox 1 c | i ma ro- | O. | | Cancer AIDS | • | | | I | |
| Wt loss in 1 month | Wt loss in 1 month Points Wt loss in 6 months | | | | | rdiac cacher | | | 1 | |
| 10% or greater 4 20% or greater | | | | | Pulmonary or cardiac cachexia 1 Presence of decubitus, open wound, or fistula 1 | | | | | |
| 5-9.9% | 3 | 10 -19.9 | | | Presence of traus | | | | ī | |
| 3-4.9% | 2 | 6-9.9 | | | Age greater than | | | | | |
| 2-2.9% | 1 | 2 - 5.9 | | | ingo areman armi | | | | | |
| 0-1-9% | 0 | 0 - 1.99 | 6 | | | | | | | |
| Score for Worksheet Record in Box | | | | | Score for Worksheet 2 = Record in Box B | | | | | |
| Worksheet 3 - Sco | ring Metabolic | Stress | | | | | | | | |
| Score for metabolic stress is | | | oles known | to increas | e protein & calorie needs. I | The score is addi | ive so that | a patieni | who has a fe | |
| | | | | | ould have an additive score | | | | | |
| Stress | none (0) | low | | | moderate (2) | | high (| | | |
| Fever | no fever | | and <10 |)1 | ≥101 and <102 | | ≥102 | | | |
| Fever duration | no fever | <72 | | | 72 hrs | | > 72 h | nrs. | | |
| Steroids | no steroids | | dose | | moderate dose | | | ose ster | rolds | |
| 41313133 | | | ing predi | nisone | (≥10 and <30mg | | | ng preda | | |
| | | | valents/ | | prednisone | | | alents/o | | |
| | | | | | equivalents/day) | | - | | حص ` | |
| | | | : | • | | Score | for Wor | ksheet 3 | · - L-4 | |
| | | | _ | | | к | ecord in | BOX C | <u> </u> | |
| Worksheet 4 - Phys | | | | | | | | | | |
| Physical exam includes a su | | | | | | | | | | |
| rated for degree of deficit. I deficit, 3+ = severe deficit. | | | | | | | | | | |
| | Kacag or deficit in me | e emellous | T MC NOI D | MOILITE OM | • | | ources (or p | TUOULUL L | A CAUCAMA I I COM | |
| Fat Stores: orbital fat pads | 10 | (1+) | 2+ | 3+ | Fluid Status: | | | | | |
| triceps skin fold | | Ψ, | 2+ | 3+ | ankie edema | (0) | 1+ | 2+ | 3+ | |
| fat overlying lower ribs | 8 | 1+ | 2+ | 3+ | sacral edema ascites | \ 0 / | 1+ 1+ | 2+ 2+ | 3+ 3+ | |
| Global fat deficit ratin | | 1+ | 2+ | 3+ | Global fluid status | \ -/ | 1+ | 2+ | 3+ | |
| Muscle Status: | | \sim | | | Point score for the phy | | | | | |
| temples (temporalis muscle | e) 0_ | (\bullet) | 2+ | 3+ | subjective rating of total | | coci il initera | <i>o</i> , | , | |
| clavicles (pectoralis & dalt | ioids) | 1+ | 2+ | 3+ | No deficit | | 0 points | | | |
| shoulders (deltoids) | 0 | (+) | 2+ | 3+ | Mild deficit | | | | | |
| interosseous muscles scapula (latissimus dorsi, trap | · · · · · · · · · · · · · · · · · · · | 1+ | 2+ | 3+ | Moderate de | ficit score = | 2 points | | | |
| | ocznas, deltoras) (0) | 4 | 2+ 2+ | 3+ 3+ | Severe defici | t score = | 3 points | | l | |
| thigh (quadriceps) calf (gastrocnemius) | ď | +(-) +(-)+(-)+ | 2+ | 3+ | | | - | | | |
| Global muscle status ri | | 1+ | 2+ | 3+ | | | or Works | | - | |
| | | | | • | | Re | cord in 1 | Box D | | |
| Worksheet 5 - PG | -SGA Global A | ssessme | nt Cate | evories | | | | | | |
| | Stage A | | سر سر | Stage | B | Stage | · C | | | |
| Category | Well-nourished | | M | | palnourished | Severely mai | | | | |
| • | | | or | suspected | malnutritico | • | | | | |
| Weight No wt loss OR | | 25 | % we loss | within I month | > 5% sur line | ia I mon | ek. | | | |
| Recent non-fluid wt gain | | (or 10% in 6 months) OR | | | > 5% wt loss in 1 month (or >10% in 6 months) OR | | | | | |
| | | | | zation or wt gain | No wt stabilization or wt gain | | | | | |
| | | | 1 | i.e., comin | ued wt loss) | | inued wt le | | | |
| Nutrient Intake No deficit OR Definite Significant recent improvement | | | una ia istaba | | | | | | | |
| | | rovement | Definite decrease in intake | | | Severe deficit | Severe deficit in intake | | | |
| | | | (B. | Presence of autrition impact Presence of nutrition impact | | | | | | |
| Nutrition Impact None OR Symptoms Significant recent improvement | | symptoms (Box 3 of PG-SGA) | | | symptoms (E | | • | | | |
| Symptoms | allowing adequate int | | 5.5 | pwnia (D | | aludaoma (E | ~ 2 01 LC | | | |
| Eventionic - | No deficit OR | nnt | | adama * | aminual deficie OR | Severe funct | lanal 3-6 | | | |
| Functioning | Significant recent im | | , | ecent deter | nctional deficit OR | recent signifi | | | | |
| | ownwert took all | to Action! | 1 4 | Sent detel | INCESTION . | terest HRIBU | vani deteli | U 18.10(1) | | |
| Physical Exam | No deficit OR | | Ev | ridence of r | mild to moderate | Obvious sign | of malnu | trition | | |
| | Chronic deficit but w | loss of SQ fat &/or muscle mass | | | (a.g., severe loss of SQ tissues, | | | | | |
| | clinical improvement | | \ & | or mpacte. | tone on paipation | possible eden | | - | | |